Vegavis

Vegavis is a <u>genus</u> of extinct <u>bird</u> that lived during the <u>Late Cretaceous</u> (<u>Maastrichtian</u> stage) of <u>Antarctica</u>, some 68 to 66 <u>mya</u>. Among modern birds, most studies show that *Vegavis* is most closely related to <u>ducks</u> and <u>geese</u> (<u>Anatidae</u>), but it is not considered to be a direct ancestor of them. [2] Although other studies question these results. [3]

Vegavis was a bird with high metabolism, which allowed it to live at high latitudes in Antarctica. It also shows a degree of <u>osteosclerosis</u>, a condition shared with <u>Polarornis</u>. This different degrees of osteosclerosis could be related to variations in diving behaviour.^[4]

Contents

Etymology

Description

See also

References

External links

Etymology

The genus name, *Vegavis*, is a combination of the name of <u>Vega Island</u> and "avis", the Latin word for bird. The species name, "iaai", is after the <u>acronym</u> for Instituto Antartico Argentino (IAA), the Argentine scientific expedition to Antarctica.

Description

The discovery of the type species, *Vegavis iaai*, demonstrates that the major groups of bird alive today had already diversified in the

Temporal range: Maastrichtian ~68-66 Ma Pre€ € OS D C P T J K PgN Life restoration based on the 2017 study by Angolín et al.[1] Scientific classification 🥖 Kingdom: Animalia Phylum: Chordata Class: Aves Order: †Vegaviiformes †Vegaviidae Family: Genus: †Vegavis Clarke et al. 2005 Species: †V. iaai **Binomial name** †Vegavis iaai Clarke et al. 2005

Vegavis

<u>Cretaceous</u>. This supports the longstanding phylogenetic inferences of <u>paleornithologists</u>. It has been hailed as the first definitive physical proof that representatives of some of the groups of modern birds lived in the Mesozoic.^[2]

The <u>holotype</u> specimen of *Vegavis* is held by the <u>Museo de La Plata</u>, <u>Argentina</u>. The specimen, cataloged as MLP 93-I-3-1, was found in the <u>López de Bertodano Formation</u> at <u>Cape Lamb</u> on <u>Vega Island</u>, Antarctica, in 1993, but was only described as a new species in 2005 because it consists of the very delicate remains of one bird embedded in a <u>concretion</u>, which had to be meticulously prepared for study. <u>CT</u> scans were utilized to gain a clearer picture of the bone structure without running danger of damaging or destroying the fossil. [2]

A second specimen, MACN-PV 19.748, was found beside the holotype specimen. It was preserved in three dimensions; CT scans were again utilized to visualize the intact <u>syrinx</u> of this specimen. The syrinx has an asymmetrical third segment, suggesting that *Vegavis* had two sources of sound in the neck and along with large soft-tissue resonating structures. This indicates that it was likely capable of honks as in ducks, geese, and other <u>basal neognaths</u>. In 2017 a phylogenetic study Agnolín and colleagues have found *Vegavis* to be stemanseriforms along with *Polarornis*, *Neogaeornis* and *Australornis* in the family Vegaviidae. [1]

See also

Asteriornis

References

- 1. Agnolín, F.L.; Egli, F.B.; Chatterjee, S.; Marsà, J.A.G (2017). "Vegaviidae, a new clade of southern diving birds that survived the K/T boundary". *The Science of Nature*. **104** (87): 87. doi:10.1007/s00114-017-1508-y (https://doi.org/10.1007%2Fs00114-017-1508-y). PMID 28988276 (https://pubmed.ncbi.nlm.nih.gov/28988276).
- 2. Clarke, J.A.; Tambussi, C.P.; Noriega, J.I.; Erickson, G.M.; Ketcham, R.A. (2005). "Definitive fossil evidence for the extant avian radiation in the Cretaceous" (http://www.digimorph.org/specimens/Vegavis_iaai/nature03150.pdf) (PDF). Nature. 433 (7023): 305–308. doi:10.1038/nature03150 (https://doi.org/10.1038%2Fnature03150). PMID 15662422 (https://pubmed.ncbi.nlm.nih.gov/15662422). Supporting information (http://www.nature.com/nature/journal/v433/n7023/suppinfo/nature03150.html)
- 3. Mayr, G. *et al.* (2018) On the taxonomic composition and phylogenetic affinities of the recently proposed clade Vegaviidae Agnolín et al., 2017 neornithine birds from the Upper Cretaceous of the Southern Hemisphere (https://www.sciencedirect.com/science/article/pii/S019566711730 5232). *Cretaceous Research*, 86:178-185.
- Marsà, Jordi Alexis Garcia; Agnolín, Federico L.; Novas, Fernando (2019). "Bone microstructure of Vegavis IAAI (Aves, Anseriformes) from the Upper Cretaceous of Vega Island, Antarctic Peninsula". *Historical Biology*. 31 (2): 163–167. doi:10.1080/08912963.2017.1348503 (https://doi.org/10.1080%2F08912963.2017.1348503).
- 5. Clarke, J.A.; Chatterjee, S.; Li, Z.; Riede, T.; Agnolin, F.; Goller, F.; Isasi, M.P.; Martinioni, D.R.; Mussel, F.J.; Novas, F.E. (2016). "Fossil evidence of the avian vocal organ from the Mesozoic". *Nature*. **538** (7626): 502–505. doi:10.1038/nature19852 (https://doi.org/10.1038%2Fnature19852). PMID 27732575 (https://pubmed.ncbi.nlm.nih.gov/27732575).

External links

- 'Dinosaurs Mingled with Cousins of Ducks and Chickens', from Live Science (http://www.livescience.com/animalworld/050125 dino chickens.html)
- 'These Are the Dinosaurs That Didn't Die', from National Geographic (https://www.nationalgeographic.com/magazine/2018/05/dinosaurs-survivors-birds-fossils/)

Retrieved from "https://en.wikipedia.org/w/index.php?title=Vegavis&oldid=946689138"

This page was last edited on 21 March 2020, at 19:56 (UTC).

Text is available under the <u>Creative Commons Attribution-ShareAlike License</u>; additional terms may apply. By using this site, you agree to the <u>Terms of Use</u> and <u>Privacy Policy</u>. Wikipedia® is a registered trademark of the <u>Wikimedia</u> Foundation, Inc., a non-profit organization.